



A space-time analysis of *Mycoplasma bovis* in Denmark

Arede, Margarida; Nielsen, Per Kantsø; Ahmed, Syed Sayeem Uddin; Hisham Beshara Halasa, Tariq; Nielsen, Liza R.; Toft, Nils

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Arede, M., Nielsen, P. K., Ahmed, S. S. U., Hisham Beshara Halasa, T., Nielsen, L. R., & Toft, N. (2015). *A space-time analysis of Mycoplasma bovis in Denmark*. Poster session presented at Annual Meeting of the Society of Veterinary Epidemiology and Preventive medicine, Ghent, Belgium.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A space-time analysis of *Mycoplasma bovis* in Denmark

Margarida Arede¹, Per K. Nielsen¹, Syed S.U. Ahmed¹, Tariq Halasa¹, Liza R. Nielsen², Nils Toft¹

¹ Section for Epidemiology, National Veterinary Institute, Technical University of Denmark

² Department of Large Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen


Background

Mycoplasma (M.) bovis causes in cattle, among other diseases, mastitis. The dairy cattle population in Denmark had an increase in atypical clinical outbreaks of *M. bovis* over the past years. An important prerequisite to the implementation of an effective control program is to **determine the geographical distribution of *M. bovis*.**

Conclusions

- *Mycoplasma bovis* infected herds are clustered in northern or southern Denmark.


Data



Results

Space-time scan statistics (SatScan™) / K- function


Significant local primary clusters*



The maps show the location of the clusters of *M. bovis* infected herds, while the inserts (K-function) indicate global clustering of cases around a radius of 70 km (app), in each round.

(*)There was no clustering in the second round.

A marginally significant primary cluster:



The clusters are in areas with high herd and cattle density.

Acknowledgements:

- Danish Milk Levy fund who funded the sampling
- SEGES who provided the data

